



Contents lists available at ScienceDirect

Journal of the Formosan Medical AssociationJournal homepage: <http://www.jfma-online.com>**News and Perspectives****Hepatitis C Infection and Metabolic Syndrome***Ching-Sheng Hsu,^{1,2} Jia-Horng Kao^{2–5*}*

Infection with the Hepatitis C virus (HCV) is a globally important cause of chronic hepatitis, cirrhosis and hepatocellular carcinoma.^{1,2} HCV infection is also associated with several extrahepatic manifestations, including mixed cryoglobulinemia, non-Hodgkin lymphoma, porphyria cutanea tarda, lichen planus, thyroid disease (usually hypothyroidism), lymphocytic sialadenitis, and polyarthritis.³ Among extrahepatic diseases, metabolic derangements of patients with chronic HCV infection have been increasingly recognized and have become an area of active investigation, especially given the global epidemic of obesity and diabetes. Although a direct link between obesity or metabolic syndrome and HCV infection remains unclear,⁴ several studies have demonstrated synergistic effects of HCV infection and obesity on the development of liver cancer.⁵ Accumulating evidence already demonstrates that HCV infection has an impact on metabolic derangements, including glucose metabolism and lipid metabolism.^{6–10} It is thus reasonable to speculate that HCV itself may elicit liver cell damage as well as metabolic abnormalities. Both events will lead to the emergence and progression of metabolic syndrome.

The interactions between HCV infection and metabolic syndrome are reviewed and discussed in this article.

HCV Infection and Glucose Metabolism

Epidemiological studies have demonstrated a higher prevalence of HCV infection in diabetic patients compared with matched controls,⁶ a higher prevalence of diabetes in HCV-infected patients,^{6,8} and an increased risk of developing diabetes in patients infected with HCV than those not infected [relative hazard = 11.58; 95% confidence interval (CI) = 1.39–96.6].¹¹ Of particular note is that a population-based survey on 10,975 Southern Taiwanese participants identified a significant link between type 2 diabetes mellitus and HCV viremia. The prevalence of type 2 diabetes mellitus in HCV viremic subjects (18.0%) was significantly higher than those in hepatitis B surface antigen (HBsAg)-positive subjects (11.4%) and subjects negative for both viral hepatitis markers (12.5%).¹² Moreover, HCV infection may induce insulin resistance (IR), and this effect seems to be genotype-specific with

©2010 Elsevier & Formosan Medical Association

¹Division of Gastroenterology, Department of Internal Medicine, Buddhist Tzu Chi General Hospital, Taipei Branch, and School of Medicine, Tzu Chi University, Hualien; ²Graduate Institute of Clinical Medicine, Departments of ³Internal Medicine and ⁴Medical Research, and ⁵Hepatitis Research Center, National Taiwan University College of Medicine and National Taiwan University Hospital, Taipei, Taiwan.

***Correspondence to:** Dr Jia-Horng Kao, Graduate Institute of Clinical Medicine, National Taiwan University College of Medicine, 7 Chung-Shan South Road, Taipei 10002, Taiwan.
E-mail: kaohj@ntu.edu.tw

a lower IR in genotype 3 than other genotypes.¹³ Our previous studies on Taiwanese chronic hepatitis C patients not only identified an association between higher HCV RNA level and a lower serum adiponectin level, a marker of IR,¹⁴ but also a dose-dependent response between homeostatic model assessment-IR index (a method of IR measurement) and HCV RNA level.¹⁵ These findings have been confirmed by a recent study.¹⁶

Several mechanisms have been proposed to explain the relationship between HCV infection, IR and diabetes, such as a direct pancreatic β cell destruction, autoimmune β cell injury, and the presence of non-alcoholic fatty liver disease.¹⁷ One study examined the liver specimens from non-obese and non-diabetic subjects with HCV infection and found that HCV infection may contribute to IR by causing an insulin signaling defect in hepatic insulin receptor substrate (IRS)-1 tyrosine phosphorylation and phosphatidylinositol 3-kinase activation.¹⁸ In addition, studies on HCV core-transgenic mice livers and HCV core-transfected human hepatoma cells demonstrated that HCV may regulate glucose metabolism by core-induced suppression of cytokine signal 3 as well as the promotion of proteasomal degradation of IRS-1 and IRS-2 through ubiquitination.¹⁹ Elevated tumor necrosis factor- α level with disturbance in tyrosine phosphorylation of IRS-1,²⁰ or expression of protein phosphatase 2A to dephosphorylate protein kinase B/Akt have also been indicated.²¹ Conversely, despite the small sequence divergence of the HCV core proteins of genotypes 3a and 1b, these two proteins appear to interfere with the insulin signaling pathway by different and genotype-specific mechanisms (Figure).²²

HCV Infection and Lipid Metabolism

Hepatic steatosis is commonly observed in subjects with chronic HCV, occurring in approximately 50% of liver biopsy samples, with a reported range of 30–70%.^{8,14,23,24} A comparison study between 317 HCV liver biopsy specimens and 299 hepatitis B viral specimens indicated that large-droplet

fat is more likely to be seen in HCV than in hepatitis B viral infection (OR=2.4; 95% CI=1.4–4.1), and Mallory body-like material is found only in HCV biopsy specimens (OR=71.6; 95% CI=4.4–996.1).²⁴ Regarding liver histologic data among patients with different HCV genotype infections, genotype 3a infection has a significantly higher prevalence of steatosis.^{9,25} In addition, the grade of hepatic steatosis is associated more with HCV RNA level in genotype 3a infection ($r=0.786$; $p<.001$), but less with body mass index in genotype 1 infection ($r=0.689$; $p<.001$).⁹ However, existing lines of evidence indicate a combination of host and viral effects on steatosis, with the relative importance of each varying with HCV genotype.^{26,27} In Taiwan, HCV genotypes 1 and 2 are more rampant than other genotypes, and hepatic steatosis in chronic hepatitis C patients was found to be associated with features of metabolic syndrome, rather than HCV genotype, advanced fibrosis or the response to antiviral therapy.¹⁰ The link between HCV and metabolic abnormalities is further supported by two population-based studies from Taiwan, one showed an association between positive anti-HCV and serum high-density lipoprotein-cholesterol level as well as serum triglyceride level,²⁸ and the other demonstrated lower serum cholesterol and triglyceride levels in HCV viremic patients.²⁹ Furthermore, our recent data identified that chronic hepatitis C patients not only have higher high-density lipoprotein-cholesterol, lower total cholesterol, triglyceride, and low-density lipoprotein-cholesterol (LDL) levels, but also higher serum adiponectin levels than matched healthy controls.⁷ Conversely, sustained disappearance of HCV is associated with reduction of steatosis in genotype 3,³⁰ and a correction of baseline low serum cholesterol and LDL.^{27,31} A recent study confirmed that HCV infection was associated with decreased cholesterol and LDL levels and this hypolipidemic effect disappeared with successful hepatitis C treatment but persisted in non-responders.³¹ Additionally, a significant portion of patients successfully treated had rebound LDL and cholesterol to levels associated with increased risk of coronary heart disease.




Figure. Hepatitis C virus affects signal transduction of insulin. The insulin receptor is tyrosine kinase-linked and undergoes auto-phosphorylation. This receptor catalyses the phosphorylation of cellular proteins such as insulin receptor substrate (IRS)-1 or 2. Upon tyrosine phosphorylation, these proteins interact with signaling molecules, resulting in a diverse series of signaling pathways, including activation of phosphatidylinositol-3-OH kinase and downstream PtdIns(3,4,5)P₃-dependent protein kinases (PIP₃-dependent protein kinases), such as phosphoinositide-dependent kinase 1 (PDK1) or atypical protein kinase C. PIP₃ interacts with and activates PDK1, which partially activates protein kinase B (PKB) by phosphorylating PKB at Thr308. Maximal activation of PKB also requires the phosphorylation of Ser473 by mammalian target of rapamycin (mTOR) complex. These pathways act in concert to coordinate the regulation of glucose, lipid and protein metabolism. Hepatitis C virus (HCV) may affect these pathways by core-induced suppression of cytokine signal, as well as the promotion of proteasomal degradation of IRS-1 and IRS-2 through ubiquitination. Alternatively, HCV may elevate tumor necrosis factor- α (TNF- α) levels disturbing tyrosine phosphorylation of IRS-1, or expression of protein phosphatase 2A (PP2A) to dephosphorylate PKB/Akt.

The mechanisms involved in the association of hepatic steatosis with HCV infection remain unclear.³² Evidence from transgenic murine models showed that expression of HCV structural proteins enhances a low background of hepatic steatosis,³³ and HCV core protein may inhibit microsomal triglyceride transfer protein activity, very low density lipoprotein secretion,³⁴ as well as alter the expression patterns of lipid metabolism genes.³⁵ Recent studies further identified the important roles of HCV core and NS5A protein on HCV-induced dyslipidemia and steatosis.^{36–39} Moreover, experiments have demonstrated the important roles of assembly and secretion of very low-density lipoproteins in HCV production,⁴⁰ and using inhibitors of lipid metabolism may suppress replication of infectious HCV clones.⁴¹ Of particular note, the lipid droplet, an organelle

for the storage of neutral lipids, has been linked to the production of infectious virus particles,⁴² which implies a linkage between HCV RNA levels and lipid profiles. We thus assessed this possible link in 531 chronic hepatitis C patients, and found a proportional relationship between serum lipid profiles and hepatitis C viral load in patients with genotype 2 infection, but not in genotype 1 infection.⁴³

Effects of Metabolic Abnormalities on HCV Infection

In addition to the impact of HCV infection on metabolic profiles, metabolic abnormalities may influence disease progression and therapeutic response in chronic hepatitis C patients. Several

studies have shown a significant and independent association between the grade of steatosis and liver fibrosis,⁴⁴ as well as IR and fibrotic progression in chronic hepatitis C.¹³ Meanwhile, obesity, defined as a body mass index greater than 30 kg/m², is a negative predictor of response to hepatitis C treatment, independent of genotype and cirrhosis.⁴⁵ Similarly, a higher IR is associated with a lower sustained virologic response in patients treated with combination therapy of peginterferon plus ribavirin,^{46,47} especially among a “difficult-to-treat” population.⁴⁸ A study evaluating 198 chronic hepatitis C patients treated with combination therapy showed that HCV genotype 1b virus may induce IR via up-regulation of suppressor of cytokine signal 3 to impair therapeutic response (OR=6.7; $p<0.005$).^{19,49} Conversely, clearance of HCV appears to improve IR, β cell function and hepatic IRS-1/2 expression.⁵⁰

Conclusions

HCV induces several metabolic abnormalities as well as interacts with the components of metabolic syndrome. These interactions vary among different HCV genotypes. In addition, each component of metabolic syndrome can directly influence the other, and thus all components should be managed simultaneously to delay or prevent disease progression and to improve therapeutic response in patients with chronic HCV infection.

References

1. Kao JH, Chen DS. Transmission of hepatitis C virus in Asia: past and present perspectives. *J Gastroenterol Hepatol* 2000;15 Suppl:E91–6.
2. Chen DS. Hepatitis C virus in chronic liver disease and hepatocellular carcinoma in Taiwan. *Princess Takamatsu Symp* 1995;25:27–32.
3. Galossi A, Guarisco R, Bellis L, et al. Extrahepatic manifestations of chronic HCV infection. *J Gastrointest Liver Dis* 2007;16:65–73.
4. Shaheen M, Echeverry D, Oblad MG, et al. Hepatitis C, metabolic syndrome, and inflammatory markers: results from the Third National Health and Nutrition Examination Survey [NHANES III]. *Diabetes Res Clin Pract* 2007;75:320–6.
5. Chen CL, Yang HI, Yang WS, et al. Metabolic factors and risk of hepatocellular carcinoma by chronic hepatitis B/C infection: a follow-up study in Taiwan. *Gastroenterology* 2008;135:111–21.
6. Mason AL, Lau JY, Hoang N, et al. Association of diabetes mellitus and chronic hepatitis C virus infection. *Hepatology* 1999;29:328–33.
7. Hsu CS, Liu CJ, Liu CH, et al. Metabolic profiles in patients with chronic hepatitis C: a case-control study. *Hepatol Int* 2008;2:250–7.
8. Mehta SH, Brancati FL, Sulkowski MS, et al. Prevalence of type 2 diabetes mellitus among persons with hepatitis C virus infection in the United States. *Ann Intern Med* 2000;133:592–9.
9. Adinolfi LE, Gambardella M, Andreana A, et al. Steatosis accelerates the progression of liver damage of chronic hepatitis C patients and correlates with specific HCV genotype and visceral obesity. *Hepatology* 2001;33:1358–64.
10. Liu CJ, Jeng YM, Chen PJ, et al. Influence of metabolic syndrome, viral genotype and antiviral therapy on superimposed fatty liver disease in chronic hepatitis C. *Antivir Ther* 2005;10:405–15.
11. Mehta SH, Brancati FL, Strathdee SA, et al. Hepatitis C virus infection and incident type 2 diabetes. *Hepatology* 2003;38:50–6.
12. Huang JF, Dai CY, Hwang SJ, et al. Hepatitis C viremia increases the association with type 2 diabetes mellitus in a hepatitis B and C endemic area: an epidemiological link with virological implication. *Am J Gastroenterol* 2007;102:1237–43.
13. Hui JM, Sud A, Farrell GC, et al. Insulin resistance is associated with chronic hepatitis C virus infection and fibrosis progression [corrected]. *Gastroenterology* 2003;125:1695–704.
14. Liu CJ, Chen PJ, Jeng YM, et al. Serum adiponectin correlates with viral characteristics but not histologic features in patients with chronic hepatitis C. *J Hepatol* 2005;43:235–42.
15. Hsu CS, Liu CJ, Liu CH, et al. High hepatitis C viral load is associated with insulin resistance in patients with chronic hepatitis C. *Liver Int* 2008;28:271–7.
16. Moucari R, Asselah T, Cazals-Hatem D, et al. Insulin resistance in chronic hepatitis C: association with genotypes 1 and 4, serum HCV RNA level, and liver fibrosis. *Gastroenterology* 2008;134:416–23.
17. Younossi ZM, McCullough AJ. Metabolic syndrome, non-alcoholic fatty liver disease and hepatitis C virus: impact on disease progression and treatment response. *Liver Int* 2009;29 Suppl 2:3–12.
18. Aytug S, Reich D, Sapiro LE, et al. Impaired IRS-1/PI3-kinase signaling in patients with HCV: a mechanism for increased prevalence of type 2 diabetes. *Hepatology* 2003;38:1384–92.
19. Kawaguchi T, Yoshida T, Harada M, et al. Hepatitis C virus down-regulates insulin receptor substrates 1 and 2 through up-regulation of suppressor of cytokine signaling 3. *Am J Pathol* 2004;165:1499–508.

20. Shintani Y, Fujie H, Miyoshi H, et al. Hepatitis C virus infection and diabetes: direct involvement of the virus in the development of insulin resistance. *Gastroenterology* 2004;126:840–8.
21. Bernsmeier C, Duong FH, Christen V, et al. Virus-induced over-expression of protein phosphatase 2A inhibits insulin signalling in chronic hepatitis C. *J Hepatol* 2008;49:429–40.
22. Paziienza V, Clement S, Pugnale P, et al. The hepatitis C virus core protein of genotypes 3a and 1b downregulates insulin receptor substrate 1 through genotype-specific mechanisms. *Hepatology* 2007;45:1164–71.
23. Scheuer PJ, Ashrafzadeh P, Sherlock S, et al. The pathology of hepatitis C. *Hepatology* 1992;15:567–71.
24. Lefkowitz JH, Schiff ER, Davis GL, et al. Pathological diagnosis of chronic hepatitis C: a multicenter comparative study with chronic hepatitis B. The Hepatitis Interventional Therapy Group. *Gastroenterology* 1993;104:595–603.
25. Rubbia-Brandt L, Quadri R, Abid K, et al. Hepatocyte steatosis is a cytopathic effect of hepatitis C virus genotype 3. *J Hepatol* 2000;33:106–15.
26. Hickman IJ, Clouston AD, Macdonald GA, et al. Effect of weight reduction on liver histology and biochemistry in patients with chronic hepatitis C. *Gut* 2002;51:89–94.
27. Poynard T, Ratzu V, McHutchison J, et al. Effect of treatment with peginterferon or interferon alfa-2b and ribavirin on steatosis in patients infected with hepatitis C. *Hepatology* 2003;38:75–85.
28. Jan CF, Chen CJ, Chiu YH, et al. A population-based study investigating the association between metabolic syndrome and hepatitis B/C infection (Keelung Community-based Integrated Screening study No. 10). *Int J Obes (Lond)* 2006;30:794–9.
29. Dai CY, Chuang WL, Ho CK, et al. Associations between hepatitis C viremia and low serum triglyceride and cholesterol levels: a community-based study. *J Hepatol* 2008;49:9–16.
30. Rubbia-Brandt L, Giostra E, Mentha G, et al. Expression of liver steatosis in hepatitis C virus infection and pattern of response to alpha-interferon. *J Hepatol* 2001;35:307.
31. Corey KE, Kane E, Munroe C, et al. Hepatitis C virus infection and its clearance alter circulating lipids: implications for long-term follow-up. *Hepatology* 2009;50:1030–7.
32. Negro F, Sanyal AJ. Hepatitis C virus, steatosis and lipid abnormalities: clinical and pathogenic data. *Liver Int* 2009; 29 Suppl 2:26–37.
33. Lerat H, Honda M, Beard MR, et al. Steatosis and liver cancer in transgenic mice expressing the structural and non-structural proteins of hepatitis C virus. *Gastroenterology* 2002;122:352–65.
34. Perlemuter G, Sabile A, Letteron P, et al. Hepatitis C virus core protein inhibits microsomal triglyceride transfer protein activity and very low density lipoprotein secretion: a model of viral-related steatosis. *FASEB J* 2002;16:185–94.
35. Chang ML, Yeh CT, Chen JC, et al. Altered expression patterns of lipid metabolism genes in an animal model of HCV core-related, nonobese, modest hepatic steatosis. *BMC Genomics* 2008;9:109.
36. Jackel-Cram C, Babiuk LA, Liu Q. Up-regulation of fatty acid synthase promoter by hepatitis C virus core protein: genotype-3a core has a stronger effect than genotype-1b core. *J Hepatol* 2007;46:999–1008.
37. Hourieux C, Patient R, Morin A, et al. The genotype 3-specific hepatitis C virus core protein residue phenylalanine 164 increases steatosis in an in vitro cellular model. *Gut* 2007;56:1302–8.
38. Moriishi K, Mochizuki R, Moriya K, et al. Critical role of PA28gamma in hepatitis C virus-associated steatogenesis and hepatocarcinogenesis. *Proc Natl Acad Sci USA* 2007; 104:1661–6.
39. Shi ST, Polyak SJ, Tu H, et al. Hepatitis C virus NS5A colocalizes with the core protein on lipid droplets and interacts with apolipoproteins. *Virology* 2002;292:198–210.
40. Huang H, Sun F, Owen DM, et al. Hepatitis C virus production by human hepatocytes dependent on assembly and secretion of very low-density lipoproteins. *Proc Natl Acad Sci USA* 2007;104:5848–53.
41. Amemiya F, Maekawa S, Itakura Y, et al. Targeting lipid metabolism in the treatment of hepatitis C virus infection. *J Infect Dis* 2008;197:361–70.
42. Miyanari Y, Atsuzawa K, Usuda N, et al. The lipid droplet is an important organelle for hepatitis C virus production. *Nat Cell Biol* 2007;9:1089–97.
43. Hsu CS, Liu CH, Liu CJ, et al. Association of lipid profiles with hepatitis C viral load in chronic hepatitis C patients with genotype 1 or 2 infection. *Am J Gastroenterol* 2009; 104:598–604.
44. Leandro G, Mangia A, Hui J, et al. Relationship between steatosis, inflammation, and fibrosis in chronic hepatitis C: a meta-analysis of individual patient data. *Gastroenterology* 2006;130:1636–42.
45. Bressler BL, Guindi M, Tomlinson G, et al. High body mass index is an independent risk factor for nonresponse to antiviral treatment in chronic hepatitis C. *Hepatology* 2003;38:639–44.
46. Romero-Gomez M, Del Mar Vioria M, Andrade RJ, et al. Insulin resistance impairs sustained response rate to peginterferon plus ribavirin in chronic hepatitis C patients. *Gastroenterology* 2005;128:636–41.
47. Conjeevaram HS, Kleiner DE, Everhart JE, et al. Race, insulin resistance and hepatic steatosis in chronic hepatitis C. *Hepatology* 2007;45:80–7.
48. Dai CY, Huang JF, Hsieh MY, et al. Insulin resistance predicts response to peginterferon-alpha/ribavirin combination therapy in chronic hepatitis C patients. *J Hepatol* 2009; 50:712–8.
49. Persico M, Capasso M, Persico E, et al. Suppressor of cytokine signaling 3 (SOCS3) expression and hepatitis C virus-related chronic hepatitis: Insulin resistance and response to antiviral therapy. *Hepatology* 2007;46:1009–15.
50. Kawaguchi T, Ide T, Taniguchi E, et al. Clearance of HCV improves insulin resistance, beta-cell function, and hepatic expression of insulin receptor substrate 1 and 2. *Am J Gastroenterol* 2007;102:570–6.